

Lepton pair production in relativistic ion collisions to all orders in $Z\alpha$ with logarithmic accuracy

S. R. Gevorkyan^{1,2} and E. A. Kuraev¹

February 1, 2008

¹ *Joint Institute for Nuclear Research, 141980, Dubna, Russia*

² *Yerevan Physics Institute, 375036, Yerevan, Armenia*

Abstract

The problem of summation of the perturbation series for the process of lepton pair production in relativistic ion collisions is investigated. We show that the amplitude of this process can be obtained in the compact analytical form, if one confines to the terms growing as a powers of logarithm energy in the cross section, the approximation of which is completely justified for such colliders as RHIC and LHC. Using this result we calculate the Coulomb corrections to the cross section of the process under consideration.

The process of lepton pair production in the collision of charged relativistic particles was investigated many years ago [1, 2] in the lowest order in the fine structure constant $\alpha = \frac{e^2}{4\pi}$. For the total cross section of the process of lepton pair production in collision of two relativistic particles with charge numbers Z_1, Z_2

$$Z_1 + Z_2 \rightarrow e^+e^- + Z_1 + Z_2, \quad (1)$$

the famous formula was obtained in 1937 by Racah [2], which in contemporary symbols reads

$$\sigma_B = \frac{28}{27} \frac{\alpha^4 (Z_1 Z_2)^2}{\pi m^2} [L^3 - 2.2L^2 + 3.84L - 1.636], \quad (2)$$

$$L = \ln(\gamma_1 \gamma_2), \gamma_i = \frac{E_i}{m_i},$$

with m electron mass, E_i, m_i the center of mass energies and masses of colliding particles.

For relativistic heavy ion collisions this cross section is huge and the problem of Coulomb corrections (higher order photon exchanges between the produced lepton pair and the Coulomb fields of colliding ions) becomes relevant. The Coulomb corrections (CC) in the case of lepton pair photoproduction off the nucleus Coulomb field was obtained in [3]. It was shown that the total cross section for the process of lepton pair photoproduction in the Coulomb field of the nucleus with charge number Z reads

$$\sigma = \frac{28}{9} \frac{\alpha^3 Z^2}{m^2} \left[\ln \frac{2E_\gamma}{m} - \frac{109}{42} - f(\alpha Z) \right], f(x) = x^2 \sum_{n=1}^{\infty} \frac{1}{n(n^2 + x^2)}. \quad (3)$$

The production of lepton pairs from the nuclear Coulomb field by virtual photons was investigated in [4]. In [5] the Bethe- Maximon result was obtained in the framework of powerful light-cone techniques [6], which allows one to trace carefully approximations done at its derivation.

As to higher order photon exchanges in the process (1), despite the intensive work done on this issue in the last years [7, 8, 9, 10, 11] the problem of allowing for CC in all orders of perturbation theory is not solved up to now. The goal of the present work is to investigate this issue based on our recent results [12] obtained for the first terms of perturbation theory for the amplitude of the process (1).

As was shown in [12], the number of remarkable cancellations between the different terms in the cross section of the process (1) takes place, which allows one to simplify the obtained expressions and may be a hint of how to solve the problem in the general case. In what follows we generalize the results of [12] to all orders of perturbation theory and using the impact factor representation get compact analytical expressions for the cross section of the process (1) true to all orders of perturbation theory (up to the constant terms).

It is convenient to separate the full amplitude of the process (1) into different terms which lead to a distinct energy dependence in the cross section, namely

$$M = M_B + M_S + M^S + M_M; \quad (4)$$

$$M^S = \sum_{n=2}^{\infty} M_n^1; M_S = \sum_{n=2}^{\infty} M_1^n; M_M = \sum_{n,m=2}^{\infty} M_n^m$$

Here M_B is the Born amplitude (see Fig.1); the terms $M_S(M^S)$ correspond to the pair production by single photon exchange between the ion with charge number $Z_1(Z_2)$ and the produced pair and any multiple photon exchanges

between the pair of leptons and the ion with $Z_2(Z_1)$ (Fig.2); the term M_M in (4) describes the cases when the pair interacts with each ion more than once (Fig.3).

We do not consider the set of Feynman diagrams (FD) accounting for the Coulomb interactions between the colliding nuclei. Such interactions were analyzed in [13], where it has been shown that interaction between ions completely canceled when the final ions are not detected.

The square of the amplitude (4) is the combination of different terms which can be classified in the cross section by their energy dependence, i.e. by different powers of big logarithm $L = \ln(\gamma_1\gamma_2)$. The Born approximation leads to the third power of L in the total cross section (see (2)). As was discussed in [12], the next powers of L come from the square of M_1^n and M_n^1 and their interference with the Born amplitude.

As to the interference $M_1^m M_n^1$, it is compensated (up to the constant) by the interference between the Born term and relevant one M_n^m , i.e. the combinations of the type $M_B M_n^m$. The same cancellations take place also in the $|M_M|^2$ and any interference of the type $M_M M_S$. Each such combination leads in the cross section to the terms growing with energy as the first power of L. To find their contribution to the cross section one must take the imaginary part of the complex logarithm $\ln(-\frac{s}{m_1 m_2} - i\epsilon) = \ln(\frac{s}{m_1 m_2}) - i\pi$ which is energy independent. Thus, if one confines to the terms in the cross section rising with energy, the square of the relevant matrix element can be cast in the following form:

$$|M|^2 = |M_B|^2 + |M_S|^2 + |M^S|^2 + 2\text{Re}(M_B^* M_S + M_B^* M^S). \quad (5)$$

Thus to calculate the cross section of the process (1) (up to the constant terms) one needs the knowledge of the Born amplitude M_B and terms relevant to single photon exchange with one of the colliding nuclei and any exchanges with the other $M_S(M^S)$.

Let us consider the main results obtained in [12] for the first terms of such types and generalize them to all orders in $Z\alpha$. The amplitude of the process (1) in the Born approximation reads [12]

$$M_B = -is \frac{(8\pi\alpha)^2 Z_1 Z_2 N_1 N_2 (1 - \beta_1)(1 - \alpha_2)}{(\vec{q}_1^2 + \beta_1^2 m_1^2)(\vec{q}_2^2 + \alpha_2^2 m_2^2)} \bar{u}(q_-) R_B v(q_+) \quad (6)$$

$$R_B = \frac{|\vec{q}_1|}{s\beta_1} [m R_1 \hat{e}_1 + \hat{e}_1 \hat{Q}_1 + 2z_+ \vec{e}_1 \vec{Q} + 2|\vec{q}_1| z_+ z_- R_1] \hat{p}_2;$$

$$R_1 = R_1(k_-, k_+) = S(k_-) - S(k_+); \vec{Q}_1 = \vec{Q}_1(k_-, k_+) = \vec{k}_- S(k_-) + \vec{k}_+ S(k_+); \quad (7)$$

$$S(k) = \frac{1}{\vec{k}^2 + \mu^2}; \mu^2 = m^2 + z_+ z_- \vec{q}_1^2; \vec{k}_\pm = \vec{q}_\pm - z_\pm \vec{q}_1; \vec{e}_1 = \frac{\vec{q}_1}{|\vec{q}_1|}; z_\pm = \frac{\beta_\pm}{\beta_1}$$

The quantities

$$N_1 = \frac{1}{s} \bar{u}(p'_1) \hat{p}_2 u(p_1); N_2 = \frac{1}{s} \bar{u}(p'_2) \hat{p}_1 u(p_2)$$

for collision of spinless ions $N_1 = N_2 = 1$ so we will omit them later on.

We use the Sudakov parameterization for the transferred momentum q_1, q_2 and the created pair momentum q_+, q_- in terms of two light-like 4-vectors $p_1^2 = p_2^2 \sim 0$ built from the momenta of colliding ions and invariant energy $s = (p_1 + p_2)^2$, and euclidean two-vectors $\vec{q}_i$ transverse to a beam direction

$$q_1 = \alpha_1 p_2 + \beta_1 p_1 + \vec{q}_1; q_2 = \alpha_2 p_2 + \beta_2 p_1 + \vec{q}_2, q_{\pm} = \alpha_{\pm} p_2 + \beta_{\pm} p_1 + \vec{q}_{\pm}.$$

For definiteness we suppose that the created pair moves in the direction of the ion with charge number Z_1 ; thus $\beta_1 = \beta_+ + \beta_-$. A similar consideration can be easily carried out for the pairs moving in the opposite direction.

Expression (6) depends on lepton transverse momenta $q_{\pm}$ through $k_{\pm}$ unlike the amplitude for the process of pair photoproduction off the Coulomb field. This is a consequence of that in our approach we project all momenta with respect to the axis of colliding ions in their centre of mass system. As to the case of photoproduction, the natural choice is the direction of incident photon. Finally, as was shown in [17, 18] the Racah formula (2) can be reproduced starting with the above expressions.

The spin structure (6) is the same in all orders of perturbation theory for the matrix elements relevant to single photon exchange with one of the ions (with any number of exchanges with another ion). This is a direct consequence of lepton helicity conservation at high energy [15, 12].

Using this property we generalize the results obtained in [12] for the first orders in α to all orders of perturbation theory and represent the amplitude relevant to single photon exchange with the ion with charge number Z_1 and with n attachments to the other (Fig.2):

$$M_n^1 = s \frac{(8\pi)^2 \alpha Z_1 (-i\alpha Z_2)^n (1 - \beta_1)}{n! (\vec{q}_1^2 + m_1^2 \beta_1^2)} \bar{u}(q_-) \Re V(q_+); \quad (8)$$

$$\begin{aligned} \Re &= \frac{|\vec{q}_1|}{s\beta_1} [mR_n \hat{e} + \hat{e} \hat{Q}_n + 2z_+ \vec{e} \vec{Q}_n + 2|\vec{q}_1| z_+ z_- R_n] \hat{p}_2 \\ R_n &= \int \frac{d^2 k_1 \dots d^2 k_n}{\pi^{n-1} \vec{k}_1^2 \dots \vec{k}_n^2} \delta^2(\vec{k}_1 + \dots + \vec{k}_n - \vec{q}_2) R_n^1(k_-, k_+, k_1, \dots k_n), \\ \vec{Q}_n &= \int \frac{d^2 k_1 \dots d^2 k_n}{\pi^{n-1} \vec{k}_1^2 \dots \vec{k}_n^2} \delta^2(\vec{k}_1 + \dots + \vec{k}_n - \vec{q}_2) \vec{Q}_n^1(k_-, k_+, k_1, \dots k_n). \end{aligned} \quad (9)$$

The quantities $R_n^1(k_-, k_+, k_1, \dots k_n) \{ \vec{Q}_n^1(k_-, k_+, k_1, \dots k_n) \}$ can be represented as the combination of $S(k)$ from (7). The expressions for the cases $n=2, 3$

are cited in [12, 14, 15]. To get the next orders one can use the recurrence relations [4]:

$$\begin{aligned} R_n^1(k_n) &= R_{n-1}^1(k_-, k_+ - k_n) - R_{n-1}^1(k_- - k_n, k_+); \\ \vec{Q}_n^1(k_n) &= \vec{Q}_{n-1}^1(k_-, k_+ - k_n) - \vec{Q}_{n-1}^1(k_- - k_n, k_+). \end{aligned} \quad (10)$$

These relations read more effectively in terms of the impact parameter representation which allows one to sum the series like $M_S(M^S)$. Introducing the Fourier transform of the functions $R_n(k_n)\{\vec{Q}_n(k_n)\}$

$$G_n^{R\{Q\}}(r_1, r_2) = \frac{1}{(2\pi)^4} \int d^2k_- d^2k_+ e^{i\vec{r}_1\vec{k}_+ + i\vec{r}_2\vec{k}_-} R_n(k_-, k_+) \{\vec{Q}_n(k_-, k_+)\}. \quad (11)$$

one immediately obtains for the relevant structures

$$\begin{aligned} G_n^R(r_1, r_2) &= \frac{1}{2(2\pi)^2} K_0(\mu|\vec{r}_1 - \vec{r}_2|) \ln^n \frac{\vec{r}_2^2}{\vec{r}_1^2}; \\ \vec{G}_n^Q(r_1, r_2) &= \frac{\mu(\vec{r}_1 - \vec{r}_2)}{2(2\pi)^2 |\vec{r}_1 - \vec{r}_2|} K_1(\mu|\vec{r}_1 - \vec{r}_2|) \ln^n \frac{\vec{r}_2^2}{\vec{r}_1^2}. \end{aligned} \quad (12)$$

Here $K_{0,1}(x)$ are the modified Bessel functions of the zero and first order. Now the summation of perturbation series can be carried out with the result

$$\begin{aligned} G_R &= \sum_{n=2}^{\infty} \frac{(-i\nu)^n}{n!} G_n^R(r_1, r_2) = \frac{K_0(\mu|\vec{r}_1 - \vec{r}_2|)}{2(2\pi)^2} \sum_{n=2}^{\infty} \frac{(-i\nu)^n}{n!} \ln^n \frac{\vec{r}_2^2}{\vec{r}_1^2} \\ &= \frac{K_0(\mu|\vec{r}_1 - \vec{r}_2|)}{2(2\pi)^2} [a^{-i\nu} - 1 + i\nu \ln a]; \end{aligned} \quad (13)$$

$$\begin{aligned} \vec{G}_Q &= \frac{\mu(\vec{r}_1 - \vec{r}_2) K_1(\mu|\vec{r}_1 - \vec{r}_2|)}{2(2\pi)^2 |\vec{r}_1 - \vec{r}_2|} [a^{-i\nu} - 1 + i\nu \ln a]; \\ \nu &= \alpha Z_2; a = \frac{\vec{r}_2^2}{\vec{r}_1^2} \end{aligned} \quad (14)$$

Applying the inverse Fourier transformation to these expressions one obtains

$$R(k_-, k_+) = \int d^2r_1 d^2r_2 e^{-i(\vec{r}_1\vec{k}_+ + \vec{r}_2\vec{k}_-)} \frac{K_0(\mu|\vec{r}_1 - \vec{r}_2|)}{2} [a^{-i\nu} - 1 + i\nu \ln a]; \quad (15)$$

$$\vec{Q}(k_-, k_+) = \int d^2r_1 d^2r_2 e^{-i(\vec{r}_1\vec{k}_+ + \vec{r}_2\vec{k}_-)} \frac{\mu(\vec{r}_1 - \vec{r}_2)}{2 |\vec{r}_1 - \vec{r}_2|} K_1(\mu|\vec{r}_1 - \vec{r}_2|) [a^{-i\nu} - 1 + i\nu \ln a]. \quad (16)$$

The squares of amplitudes entering (5) are determined by these two structures. Really, as was shown in [12],

$$\frac{1}{4s^2} \sum |\bar{u}(q_-) \Re v(q_+)|^2 = (17)$$

$$\frac{1}{2} z_+ z_- \vec{q}_1^2 \left\{ (m^2 + 4z_+^2 z_-^2 \vec{q}_1^2) R^2 + (z_+^2 + z_-^2) \vec{Q}^2 + 4z_+ z_- (z_+ - z_-) R \vec{q}_1 \vec{Q} \right\}.$$

As to the amplitude M_S it can be obtained from M^S by simple replacements [12]

$$\vec{q}_1 \rightarrow \vec{q}_2; \vec{e}_1 \rightarrow \vec{e}_2; p_1 \rightarrow p_2; \beta_1 \rightarrow \alpha_2.$$

The differential cross section of the process (1) is connected with the full amplitude (4) by the relation

$$d\sigma = \frac{1}{8s} \sum |M|^2 d\Gamma. \quad (18)$$

The summation is carried over the final particles polarization. The phase space volume for four particles in the final state can be expressed through the variables introduced above and phase volume of the final leptons

$$d\Gamma_{\pm} = \frac{d^3 q_+}{2E_+} \frac{d^3 q_-}{2E_-} \delta^4(q_1 + q_2 - q_+ - q_-) \quad (19)$$

in the following way

$$d\Gamma = \frac{1}{(2\pi)^8} \frac{d^3 p'_1}{2E'_1} \frac{d^3 p'_2}{2E'_2} \frac{d^3 q_-}{2E_-} \frac{d^3 q_+}{2E_+} \delta^4(p_1 + p_2 - p'_1 - p'_2 - q_+ - q_-) \quad (20)$$

$$= \frac{ds_p}{4(2\pi)^8 s(1 - \alpha_2)(1 - \beta_1)} \frac{d\beta_1}{\beta_1} d^2 \vec{q}_1 d^2 \vec{q}_2 d\Gamma_{\pm}; s_p = (q_+ + q_-)^2.$$

The expressions (5-8, 15-20) allows one to calculate the fully differential cross section of the process under consideration.

Let us consider the cross section integrated over transverse momenta of final leptons. For this quantity one can obtain the simple analytical formula from which the special cases can be easily deduce.

First of all let us note that the integration over created particles transverse momenta in the combinations of type $|M_S|^2 + 2\text{Re} M_S M_B^*$ leads to the

following structures:

$$J_R = \int d^2r_1 d^2r_2 K_0^2(\mu | \vec{r}_1 - \vec{r}_2 |) [(a^{-i\nu} - 1 + i\nu \ln a)(a^{i\nu} - 1 - i\nu \ln a) + i\nu \ln a(a^{-i\nu} - 1 + i\nu \ln a) - i\nu \ln a(a^{i\nu} - 1 - i\nu \ln a)] \quad (21)$$

$$= \int d^2r_1 d^2r_2 K_0^2(\mu | \vec{r}_1 - \vec{r}_2 |) [2 - a^{i\nu} - a^{-i\nu} - \nu^2 \ln^2 a];$$

$$J_Q = \int d^2r_1 d^2r_2 K_1^2(\mu | \vec{r}_1 - \vec{r}_2 |) [2 - a^{i\nu} - a^{-i\nu} - \nu^2 \ln^2 a]. \quad (22)$$

The further integration leads to the relations [4]

$$J_R = -f(\nu) \frac{8\pi^2 \nu^2}{3\mu^4}; \quad (23)$$

$$J_Q = -f(\nu) \frac{16\pi^2 \nu^2}{3\mu^4},$$

where the function $f(\nu)$ is given by (3).

Using the above expressions and representing the cross section as the sum of Born term and the term relevant to CC contribution $d\sigma = d\sigma_B - d\sigma_C$ we obtain the following formula:

$$d\sigma_C = \frac{4\nu_1^2 \nu_2^2}{3\pi m^2} (f(\nu_1) + f(\nu_2)) \quad (24)$$

$$\times \frac{[3 + 4z(1-z)(\frac{q^2}{2m^2} - 1)] (1-\beta)d\beta}{(1+z(1-z)\frac{q^2}{m^2})^2} \frac{xdx}{\beta} \frac{1}{(x+\beta^2)^2} dz.$$

For the sake of simplicity we consider the case of ions with equal mass $m_1 = m_2 = M$ and consequently the same Lorentz factors $\gamma_1 = \gamma_2 = \gamma = \frac{E}{M}$. The variables $x = \frac{\vec{q}^2}{M^2}$ and β belongs to the intervals

$$0 \leq x < \infty; \quad (25)$$

$$\frac{m}{M\gamma^2} (1 + \frac{M}{4m}x) \leq \beta \leq 1.$$

The expression (24) becomes invalid in the case of pair production by the very soft photons ($\omega \sim 2m$ in the rest system of one of the ion) which corresponds to the lower limit of β (we are working in the center of mass system of colliding ions.) At such low energies one has to take into account the threshold effects and deviation from the eikonal approach [19] which is out of the present work scope.

The integrations in (24) can be carried out (for details see Appendix). We

cite here the expression for CC contribution to the total cross section of the process (1)

$$\sigma_C = \frac{28\nu_1^2\nu_2^2}{9\pi m^2}(f(\nu_1) + f(\nu_2))[\ln^2(\gamma_1\gamma_2) + \frac{20}{21}\ln(\gamma_1\gamma_2) + O(const)]. \quad (26)$$

The leading term in this formula $\sim \ln^2(\gamma_1\gamma_2)$ coincides with the relevant one found in [10, 11]. As to the next to leading term ($\sim \ln(\gamma_1\gamma_2)$) in (26) it differs from relevant one in [11] (see their formula (17)). The reason is transparent. We calculated next to leading terms in the part of the cross section corresponding to the single photon exchange from one of the nuclei (M_S, M^S). As to the formula (17) in [11] by derivation it is relevant to the part of the cross section provided by two or more exchanges from both nuclei (M_M with the preceding notation). As was discussed above and in [12] such exchanges contribute only to the constant part of the total cross section and are outside of our consideration. There is correspondence between (26) and relevant formula (30) in [16] obtained for the total cross section of e^+e^- pair production in the Coulomb field of the nucleus by high energy muons using the generalized Weizsacker-Williams approximation.

In deducing (26) we integrate over β from its lower limit, which is strictly speaking incorrect by the reasons mentioned above. Nevertheless the error in such approach can not be crucial and expression (26) can be safely used for the simple estimates of the Coulomb corrections to the total cross section of the process under consideration. The reason is relative smallness of the near threshold contribution of the pair photoproduction in the Coulomb field [19]. The obtained in the present work expressions can be used for calculation of completely (four particles in the final state) or partly (see (24)) differential cross sections the issue which is necessary in any experiment.

Acknowledgments

We are grateful to V.Serbo and A.Tarasov for useful discussions and interest to the work. We acknowledge the warm hospitality at ECT ,Trento.(E.K) is grateful to the Institute of Physics SAS (Bratislava), where part of this work was done. The work is supported by INTAS grants 30494,00366.

Appendix A

Let us calculate the total cross section of the lepton pair production off the nuclear Coulomb field by virtual photons i.e. the process

$$\gamma^*(k) + Z \rightarrow e^+(q_+) + e^-(q_-) + Z \quad (\text{A.1})$$

This allow us to compare the results obtained in our approach with the well known one and show how the integration over phase space can be carried out. For our purposes it is sufficient to calculate the total cross section of this process in the Born approximation. The generalization to all orders in α using our approach is an easy task.

Using the same approximations as before, the total cross section of the process (A.1) in the Born approximation read

$$d\sigma = \frac{2\alpha^3 Z^2}{\pi^2(\vec{q}^2 + q_L^2)^2} [(m^2 + 4z^2(1-z)^2 Q^2) R_S^2 + (z^2 + (1-z)^2) \vec{R}_V^2] dz d^2 q_+ d^2 q; \quad (\text{A.2})$$

$$R_S = \frac{1}{\vec{q}_+^2 + \mu^2} - \frac{1}{\vec{q}_-^2 + \mu^2}; \quad \vec{R}_V = \frac{\vec{q}_+}{\vec{q}_+^2 + \mu^2} - \frac{\vec{q}_-}{\vec{q}_-^2 + \mu^2}; \quad (\text{A.3})$$

$$\vec{q} = \vec{q}_+ + \vec{q}_-; \quad \mu^2 = m^2 + z(1-z)Q^2; \quad q_L = \frac{\mu^2 + \vec{q}_+^2}{2z(1-z)\omega}.$$

Here Q^2, ω are the mass and energy of virtual photons. The integration over transverse momenta q, q_+ can be done analytically. We divide the integration over the two-dimensional transfer momentum q in the two regions: 1) Small momentum transfer where the longitudinal momentum q_L has to be taken into account $0 \leq q \leq \xi \sim m$ and 2) the region where one can safely ignore it, i.e., $\xi \leq q \leq \infty$.

Using this trick the integration over transverse momenta $\vec{q}, \vec{q}_+$ can be carried out. As a result, the total cross section for the lepton pair production by transverse and longitudinal photons in the Coulomb field can be represented as

$$\sigma_T = \frac{4\alpha^3 Z^2}{3} \int \left[\ln \frac{2\omega z(1-z)}{\mu} - \frac{1}{2} \right] \left[\left(\frac{m}{\mu} \right)^2 + 2(z^2 + (1-z)^2) \right] \frac{dz}{\mu^2}; \quad (\text{A.4})$$

$$\sigma_S = \frac{16\alpha^3 Z^2}{3} Q^2 \int z^2(1-z)^2 \left[\ln \frac{2\omega z(1-z)}{\mu^2} - \frac{1}{2} \right] \frac{dz}{\mu^4}. \quad (\text{A.5})$$

The total cross section for the pair production by transversally polarized photons σ_T turns out to the Bethe-Maximon result (3) at $Q^2=0$. Moreover, the expression (A.4) is in agreement with (45) from [4]. As to the pair production by longitudinal photons expressions (46,59) in [4] have a misprint: the

combination $z(1-z)$ enters them in the first power in contradiction with our result (A.5).

References

- [1] L.D.Landau and E.M.Lifshitz,Phys.Z.Sowjet.6 (1934) 244
- [2] G. Racah, Nuovo Cim. 14 (1937) 93.
- [3] H. A. Bethe and L. Maximon, Phys. Rev. 93 (1954) 788.
- [4] D.Yu. Ivanov and K. Melnikov, Phys. Rev. D57 (1998) 4025.
- [5] B.Z.Kopeliovich,A.V.Tarasov,O.O.Voskresenskaya,Eur.Phys.J.A11 (2001) 345
- [6] J.M.Bjorken,J.B.Kogut,D.E.Soper,Phys.Rev. D3 (1971) 1382
- [7] A.J.Baltz and L.McLerran,Phys.Rev. C 58 (1998) 1679
- [8] A.J.Baltz,F.Gelis,L.McLerran,A.Peshier,Nucl.Phys.A695 (2001) 395
- [9] U.Eichmann,J.Reinhardt,S.Schramm,W.Greiner,Phys.Rev.A59 (1999) 1223
- [10] D.Yu. Ivanov, A. Schiller, V.G. Serbo,Phys.Lett.B454 (1999) 155.
- [11] R.N.Lee and A.I.Milstein,Phys.Rev. A64 (2001) 032106
- [12] E. Bartoš, S. R. Gevorkyan, E. A. Kuraev and N.N.Nikolaev, Phys.Rev.A66 (2002) 042720
- [13] E.Bartoš,S.R.Gevorkyan,E.A.Kuraev and N.N.Nikolaev, Phys. Lett. B538 (2002) 45.
- [14] I.F.Ginzburg,D.Yu.Ivanov,Nucl.Phys. B388 (1998) 376.
- [15] I.F.Ginzburg,S.L.Panfil and V.G.Serbo, Nucl.Phys.B284 (1987) 685.
- [16] D.Yu.Ivanov,E. A. Kuraev, V. G. Serbo and A. Schiller, Phys.Lett. B 442 (1998) 453
- [17] E. A. Kuraev and L. N. Lipatov, Yad. Fiz. 16 (1972) 1060.
- [18] V.M.Budnev,I.F.Ginzburg,G.V.Meledin,V.G.Serbo,Phys.Rep.C15 (1975) 181

- [19] A.I.Akhiezer,V.B.Berestetski,Quantum Electrodynamics,
Nauka,Moscow,1981




Figure 1: The lowest order Feynman diagrams (Born approximation)




Figure 2: The diagrams relevant to the terms with one single exchange $M_S(M^S)$.




Figure 3: Some diagrams describing the amplitudes with multiple exchanges M_M .